



PLATEROT
AESTHETICS

Sculptra Aesthetic Consent to Treat Form

I have requested that Dr. Carmelo Plateroti, D.O., Dr. Allison Pontius MD, Nikkole Souza, R.N., or Diane Powell, R.N. use Sculptra Aesthetic to attempt to restore a natural, and youthful look. Sculptra Aesthetic® is an injectable implant that contains microparticles of L-Poly-Lactic Acid, a biocompatible, biodegradable synthetic polymer from the alpha hydroxy acid family. A treatment regimen of SCULPTRA consists of up to 4 injection sessions scheduled 3-4 weeks apart.

SCULPTRA may provide cosmetic correction of nasolabial fold contour deficiencies and other facial wrinkles, including the correction of fine lines and wrinkles in the cheek region.

1. I understand that I should not use this product if I am allergic to subcutaneous, absorbable suture material.
2. I understand that the injection procedure reactions are: bruising, hematoma formation, discomfort, and erythema.
3. I understand that the risks of these injection procedure reactions are greater if I have been drinking any alcohol, taking any aspirin (Advil, Motrin, Ibuprofen, Naproxen, Aleve, etc.), or taking large doses of Ginkgo Biloba or more than 400 IU of Vitamin E per day.
4. I understand that the most common device-related adverse effect is the delayed occurrence of subcutaneous papules, which were confined to the injection site and were typically palpable (could be felt), asymptomatic and non-visible.
5. I understand that it takes several weeks to see the effects of this treatment, and the appropriate way to do this is to "treat, wait and reassess" in 4-6 weeks.
6. I understand that several vials of Sculptra Aesthetic® may be necessary to achieve the results I desire.
7. I understand that Sculptra Aesthetic® should not be injected while you have an active skin infection or inflammation in the treatment area and should not be injected into the red area of the lip. Use in the skin near the eyes is not recommended.
8. I understand the side effects may include injection site discomfort, redness, bruising, bleeding, itching and swelling. Other side effects may include small lumps under the skin that are sometimes noticeable when pressing on the treated area. Larger lumps, some with delayed onset with or without inflammation or skin discoloration. In a clinical study, the numbers of small and large lumps were low and most resolved without treatment.

I have been advised of the risks involved in such treatments, the expected benefits of such treatment, and the alternative treatments, including no treatment at all. The Physician has also indicated that with any procedure there is always the possibility of an unexpected complication.

Signature: _____ Date: _____

Printed name: _____